

Lessons Learned During Adherence Club Implementation

The Witkoppen Experience



JOHNS HOPKINS
BLOOMBERG
SCHOOL of PUBLIC HEALTH



THE UNIVERSITY
of NORTH CAROLINA
at CHAPEL HILL



USAID
FROM THE AMERICAN PEOPLE



Adherence Clubs

- RCT comparing RIC and viral suppression of clinic-based vs. community based clubs
- 12 pairs (24 clubs), 770 participants
- Patients randomized into clinic or community clubs
- Acceptability study nested with main study
 - Acceptability questionnaire for all at 12 & 24 months
 - In-depth interviews with selected participants



Implementation – General Considerations

- Facility readiness
 - Buy in from facility staff
 - Infrastructure: Adequate filing, booking systems; space; scripting, prepacking and dispensing capacity
- Recruitment efforts
 - Structured recruitment process to ensure eligible patients synchronized into clubs



Community Based Clubs - Implementation

- Community stakeholders buy in
 - Diepsloot stakeholder meeting
 - CAF meeting
- Venue
 - Selection criteria – spacious, easily accessible, multi-purpose, free
 - Within walking distance or easily accessible with public transport
 - Venues include church, multi-purpose hall, DoSD halls, NGO facilities
 - MOUs with all partners
 - All venues work well currently

Community-Based Clubs - Challenges


- Transport of medication, personnel, blood (cold chain)
 - Facilitator with driver's license
 - Car
 - Visits structured such that bloods drawn at end of club
- Pre-packed medication – avoiding distribution of incorrect medication
 - 2 tier system of checking correct medication prepacked and delivered
- Referrals of positive medical screens
 - Facilitator drives cases needing immediate attention to clinic
 - Other cases referred
- Annual medical visits
 - Performed at facility



USAID
FROM THE AMERICAN PEOPLE



Acceptability Sub-study (590 Questionnaires)



USAID
FROM THE AMERICAN PEOPLE



Conclusions

- Clubs preferred alternative to standard clinic care
- Satisfaction is similar by club location
- Concerns over lack of privacy, HIV disclosure and personal safety were low
- Community-based clubs face more logistical challenges
- Planning and design should be individualised and flexible
- Further evidence on clinical outcomes by adherence club location are needed to inform implementation and scale-up



This presentation is made possible by the generous support of the American people through the United States Agency for International Development (USAID). The contents are the sole responsibility of Witkoppen Health and Welfare Centre and do not necessarily reflect the views of USAID or the United States Government

